

# **Community Health Workers: a review of concepts, practice and policy concerns<sup>1</sup>**

Prasad BM\*  
VR Muraleedharan\*\*

August 2007

\*Prasad BM, BDS, MPH, Project Officer, CREHS, IIT Madras, Chennai, India

\*\*VR Muraleedharan, PhD, Professor of Economics, Department of Humanities and Social Sciences, IIT Madras, Chennai, India

<sup>1</sup> This review is a part of ongoing research of International Consortium for Research on Equitable Health Systems (CREHS), funded by UK Government Department for International Development (DFID) lead by London School of Hygiene & Tropical Medicine (LSHTM), UK. For more details please visit <http://www.crehs.lshtm.ac.uk/>

## 1. Introduction:

The global policy of providing primary level care was initiated with the declaration of Alma-Ata in 1978s. The countries signatory to Alma Ata declaration considered the establishment of CHW program as synonym with Primary Health Care approach (Mburu, 1994; Sringerinyuang, Hongvivatana, & Pradabmuk, 1995). Thus in many developing countries PHC approach was seen as a mass production activity for training CHWs in 1980s (Matomora, 1989). During these processes the voluntary health workers or CHWs were identified as the third workforce of “Human resource for Health”<sup>1</sup>(Sein, 2006 ). Following this approach CHWs introduced to provide PHC in 1980s are still providing care in the remote and inaccessible parts of the world (WHO, 2006a).

In this paper we attempt to (a) provide an overview of the concepts and practice of Community Health Workers (CHWs) from across a range of (developing and developed) countries, and (b) draw some insights into policy challenges that remain in designing effective CHW schemes, particularly in the Indian context. In the subsequent sections, we provide a review of the various ways in which community health workers have been deployed in different settings. To arrive at this we adopted a systematic search of literature on CHWs, using key words such as community health worker, primary health care worker, community based health care worker, lay health worker, we also used the inclusion criteria that WHO adopted for describing CHWs (WHO, 2006a), in Pub-Med, Science Direct, WHO and World Bank sources. A total of 110 studies (including Journal articles, Reports etc are mentioned in the tables) were identified for this purpose. We have classified these into three parts, namely those related to (1) design and role of CHWs (Table 1), (2) management of CHWs (Table 2), and (3) factors influencing performance of CHWs (Table 3, 4 and 5). As the reader will notice, these issues overlap and some studies refer to all three issues while most others primarily cover one of these issues. We propose this classification for reviewing the literature for analytical purpose. While our review draws upon these studies, we have indicated only a portion of them in the text.

---

<sup>1</sup> “Human Resources for Health” (HRH) is defined as the stock of all individuals engaged in improvement of health of population. They include professionals (doctors, nurses, pharmacists, lab technicians etc), non-professionals (auxiliary midwives, health visitors, dais, etc) they may be regulated or unregulated, voluntary care givers (voluntary Dots provider) and family members (JLI, 2004).

## 2. CHWs: an overview of concepts and practice

The CHWs have evolved with community based healthcare programme and have been strengthened by the PHC approach. However, the conception and practice of CHWs have varied enormously across countries, conditioned by their aspirations and economic capacity. This review identified seven critical factors that influence the overall performance of CHWs which are discussed in this section. In discussing these issues, our aim is to (a) highlight certain empirical knowledge and (b) point out, if any, gaps in the design, implementation and performance of CHWs.

1. **Gender:** Most countries have largely relied on females as CHWs (Table. 1). Although both men and women are employed at grass-roots level, there is a collective impression (particularly amongst policy makers) that female workers are able to deliver care more effectively than male workers at community level. While this may be true of maternal and child health (MCH) related services, the role of male workers in the control of epidemics (in the past) such as cholera, small-pox, plague, at the community level has been substantial across countries.<sup>2</sup> However, there has been an explicit policy-shift in India to replace male health workers by female workers at community level (GOI, 1997).
2. **Selection of CHWs:** Most studies highlight the need for recruiting CHWs from communities they serve, but they also point out the difficulties in implementing this approach<sup>3</sup>. CHWs are from the communities they serve presumably will not only be more accessible but also be able to gain the confidence of community members (Ruebush, Weller, & Klein, 1994). Experiences have shown that CHWs recruited from local communities have had greater impact on utilization, creating health awareness and health outcomes (Bang et al., 1994; Abbatt, 2005; Lewin, Dick, Pond, Zwarenstein, Aja, Wyk et al., 2005) (for example in India, AWARE in Andhra Pradesh, CINI in Kolkata, CRHP in Jamked, RUHSA in Tamil Nadu, and SEARCH in Maharashtra (Antia & Bhatia, 1993). Pakistan

---

<sup>2</sup> Impression drawn from interview with various officials in India

<sup>3</sup> For example, the social and economic class and caste background of CHWs may influence their acceptance by members of the community they serve , (Jobert, 1985)

(OPM, 2002; Douthwaite & Ward, 2005) refer table 1, sl no. 10), China (Campos, Ferreira, Souza, & Aguiar, 2004) refer table 1, sl no. 19).

3. **Nature of employment, Career prospects and Incentives:** Many studies have highlighted the role of nature of employment, career prospects and other incentives in determining the overall performance of community workers (Ballester, 2005). The experience is quite varied in the employment of CHWs across countries. In several countries, particularly in government health systems, CHWs were employed on voluntary basis and on full-time basis (refer Table 1). There are also countries that employed CHWs on contract or as regular employment with a fixed monthly salary paid by the government, such as in India (GOI, 1956). But India also has had the experience of having community health workers on voluntary basis (during 80s particularly) in the public sector (Lesile, 1985). While the experience of NGOs is also quite varied in this respect, we can safely state that there is perhaps more display of voluntarism in this sector in under-served areas (Antia & Bhatia, 1993).<sup>4</sup> The critical question that comes through the review is that not only would payment or voluntarism per se influence CHWs' performance, but its influence also depends on other factors inter alia highlighted here (Table 2 and 5).
4. **Educational Status:** The review shows that in most countries CHWs have had education up to primary level education, with 8 to 10 years of schooling (Table 1). Studies have shown that CHWs with higher educational qualifications have opportunities for alternative employment and therefore migrate from one job to another (Brown, Malca, Zumaran, & Miranda, 2006) refer table 5, sl no. 8). On the other hand it has also been highlighted that those with higher education could learn and enhance their skill in the diagnosis of common illness (Ande, Oladepo, & Brieger, 2004; Bentley, 1989) and thereby deliver better care to the community. Experience from other regions namely in Uganda shows that factors like

---

<sup>4</sup> Conclusions drawn from interviews with various NGOs on their role in the revised national tuberculosis control programme

age, sex, education and number of offspring was inconsequential on ability to classify Pneumonia and provide treatment accordingly by the CHWs (Kallander, Tomson, Nsabagasani, Sabiiti, Pariyo, & Peterson, 2006).

5. **Population and service coverage:** Two inter-related critical questions being faced at grass-roots level are: (a) “What is the optimal population size that a CHW could cover and (b) What is the optimal range of services that a CHW could deliver?” Experience across countries varies (Table 2). There are countries such as Sri Lanka where a CHW covers as low as 10 households offering a set of MCH related services ((UNICEF, 2004) refer table 1, sl no. 14). On the other hand, there are countries such as India, where a CHW covers about 1000 households (approximately 5000 populations, usually spread over 5 to 10 villages, refer table 1, sl no. 39) (UNICEF, 2004). In most countries, CHWs offer more preventive services than curative services (Salmen, 2002) (Table 2). Studies have also shown that such an approach may have reduced the confidence of the community on the effectiveness of CHWs (Bentley, 1989; Menon, 1991). CHWs in India offer a wider range of services through CHWs. The rationale for this is that it is necessary to integrate a range of services at community level in order to have better health outcomes (Table 3). But such an approach has also led to criticisms from various quarters that it has increased the overall work-load of CHWs and thereby reducing their performance (SARDI).
6. **Training:** The aspect of induction and continuing training programmes for CHWs have received considerable attention, as they are often selected without any prior experience or professional training in community health (Abbatt, 2005). In Nicaragua in 1980s CHWs were as young as 15 years old and were given a short duration training (not longer than 2 weeks, (Bender & Pitkin, 1987) refer table 2, sl no.6) particularly in curative services. These were exceptions necessitated by the political turmoil of that period in such countries. Despite such exceptions, CHWs in countries such as India receive training for about 3 months, while in other countries as such Brazil they receive training for about 6 to 8 months at the

beginning of their career (Campos et al., 2004; Leslie, 1985)( refer table 2, sl no 11 and 23). Career prospects for CHWs and their aspirations do influence their performance. For example some studies from the United States of America (Ballester, 2005; Scott & Wilson, 2006) have shown a significant drop out of CHWs due to lack of career prospects. Thus career prospects along with salaries are strong incentives in not only retaining CHWs, but also in enhancing their performance. The empirical analysis on the contents and approach of various training programs and their influence on performance of CHWs have been minimal. For example the algorithm developed by WHO on managing multiple childhood illness was found to be ineffective as CHWs reported serious difficulties in understanding training manuals(Kelly, Osamba, & Grag, 2001) and similar findings were reported in India by a Oxfam study about CHWs having difficulty in understanding training manuals(Ramprasad, 1988). The findings from the national survey on CHWs in the US suggest on the job-training to overcome these difficulties in understanding training manual (Kash, May, & Tai-Seale, 2007).

7. **Feedback, monitoring mechanisms and community participation:** Referrals and records-keeping are often highlighted for establishing a good monitoring system (Jerden, Hillervik, Hansson, Flacking, & Weinehall, 2006). Nevertheless only a few studies have brought out the importance of building healthy “inter-relationships” and “trust” among health professionals in building an effective feedback and referral systems in place [(Bhattacharyya, Winch, LeBan, & Tien, 2001) and refer table 4] For example, a study in South Africa describes the relationships between professional nurses and CHWs and how one viewed the other as a “threat” in their career (Doherty & Coetzee, 2005) refer table 4, sl no. 18). We argue that in such unhealthy competitive situations it is not possible to have an effective “referral system” in place (May & Contreras, 2006). However, the Namibian experience shows that through mutual understanding on agreed roles and responsibilities it would be possible to

have positive inter-personal relationship (Low A. & Ithindi, 2003). Studies for example in Columbia, have also shown that “feedback and rewards from the community” are more significant in the overall motivation and performance of CHWs (Robinson & Larsen, 1990) refer table 5, sl no. 2). The critical issues that still remain in this respect are: (a) How does a feedback mechanism from the community work? (b) What kinds of rewards are expected of the CHWs from the community? (c) How do they reflect the degree of trust and confidence that CHWs have gained from the community? (Arole, 2007).

### **3. Policy Challenges in design of CHW programme.**

The above review highlights several aspects to be kept in mind in designing and implementing effective CHW schemes. The review emphatically shows that (a) the selection of CHWs from the communities that they serve and (b) population-coverage and the range of services offered at the community levels are vital in the design of effective CHW schemes. It should be noted that smaller the population coverage, the more integrated and intensive the service offered by the CHWs.

The extent to which other factors should be taken into account is contingent on local conditions including the economic and socio-political factors. While the review has highlighted the role of gender, education, training, feedback and monitoring system, and incentives and career prospects, economic resource base and political commitment will largely determine the amount of attention they receive in the design and implementation of CHW schemes (Haines et al., 2007). For example, while it is obvious that good training is essential for CHWs, the contents and duration of training could be decided only along with decision on the range and nature of services to be offered by them, and the level of education that they already possess. It has been highlighted that in general there has been a lack of performance due to inadequate capacity of training institutions and lack of capacity of trainers to understand the local community structure (Global HealthTrust, 2003). Studies have shown that many CHW schemes do not provide primary curative care. Hence care should be taken while deciding the range and nature of services

that CHWs should provide in a given population. It is essential to strike a balance between preventive and curative services to be provided by them. Likewise, the role of incentives and career prospects should proceed from other design elements, such as the overall work-load (in term of population coverage, and services offered and the degree of follow up required by the CHWs) (Ofosu-Amaah, 1983). In this process, the degree of voluntarism that prevails among community members will also influence the extent to which financial incentives and career prospects need attention in the design of CHWs. It has been brought out in Doulas community health care programme-based study in North America, in where more than half of the CHWs were looking forward to be a qualified health professional preferably a nurse (Low, L. K., Moffat, A., & Brennan, P., 2006).

We measured the overall performance of CHWs that may determine the enthusiasm and motivation and continuity of the CHW schemes (Stock-Iwamoto & Korte, 1993). Often performance is measured in terms of improvement in health status of the population that CHWs serve, increase in the utilization of services provided by them, reduction in the wastage of resources, the presence and accessibility of CHWs to the community members, etc (Table 3). Computing each of these measures is data intensive and also requires careful effort in documentation and analysis over a period of time. However what is eventually important in sustaining the motivation of CHWs to function with commitment and effectiveness, as the experimentation in Parinche (FRCH-PUNE Project) (Antia & Bhatia, 1993) and SEARCH (Gadchiroli, Maharashtra) (Bang et. al., 1994) (Gryboski, Yinger, Dios, Worley, & Fikree, 2006) is the degree of trust and confidence of the community members that CHWs have gained over a period of time.

Table 6 summarizes our version of the strengths, weaknesses, opportunities and threats in the concept of CHWs from the literature we have reviewed. Such a classification of role of CHWs may have some pedagogic value. Our review shows that the whatever evidence that we already have lends support to the view that a carefully designed and implemented community health workers scheme could have far reaching implications for the whole society beyond generating better health outcomes(WHO, 1989). For example, it could improve their self-esteem (Roman, Lindsay, Moore, &



Shoemaker, 1999) refer table 4, sl no. 12), substantially empower women from low-income countries (Sundararaman, 2007) (Kovach & Worley, 2004) refer table 3, sl no. 8), and help them to earn respect from the community (Brown et al., 2006; Swider, 2002) table 6). Thus a well designed and implemented CHW scheme could help reduce social inequity.

Annexure:

**Table: 1. Profile of CHWs across different Countries**

SI no.	Author	Country	Year	Name	Age	Gender	Coverage	Empl oy*	Level of Education
1.	(Lehmann, Friedman, & Sanders, 2004)	Ghana Nigeria Kenya Tanzania Somalia	1970 1974	VHW	20-45	M:F F  M	-	FT	Literate Primary Schooling
2.	(Hathirat, 1983)	Thailand	1979	CHW	Varied	Male	--	FT	Graduates
3.	(Couper, 2004)	Iran	1979	Behvarz	Varied	M/F	1200-1600 indi	FT	Secondary graduates
4.	(Scholl, 1985)	Nicaragua	1981	Brigadista	15-19	F 55.5% M 45.5%	-	-	-
5.	(Bender & Pitkin, 1987)	Costa Rica Nicaragua Colombia	-	RHA Brigadista HP	- 13-40 -	M/F M/F M/F	1/400 HHS - 1/3000 to 4000 indi	FT	-
6.	(Reis, Elder, Satoto, kodyat, & Plamer, 1991)	Indonesia	1990	Kader	20-40	-	1/100 indi	FT	Educated
7.	(Nyonor, Awoonor-Williams, Phillips, Jones, & Miller, 2005)	Ghana	1990	VHW	-	-	3000 indi	FT	-
8.	(Ruebush et al., 1994)	Guatemala	-	CVs	12-76	M	1/100 indi	FT	-
9.	(Perez, Findley, Mejia, & Martinez, 2006)	USA	2000	CHW	20-29	F	300 indi	FT	High school
10.	(Oxford Policy Management, 2002),(Douthwait e & Ward, 2005)	Pakistan	2002	LHW	29 mean	F	1000 indi	FT	50% metric
11.	Algeria (WHO, 2006b)	Algeria	2002	CHW	-	-	-	-	-
12.	(UNICEF, 2004)	Nepal	2003	FCHV	>20	F	1/400; 1/250; 1/150 indi	FT	educated
13.	(UNICEF, 2004)	Bhutan	2003	VHW	-	M/F	20 -30 HHS	FT	--
14.	(UNICEF, 2004)	Sri Lanka	2004	CHW		M/F	1/10 HHS		Educated
15.	(Magongo, 2004)	Gautang	2004	CHW	-	-	200 HHS	FT	-
16.	(Friedman, 2005)	South Africa	2004	CHW	-	-	80 to 100 rural & 100- 150 urban HHS	FT	-
17.	(UNICEF, 2004)	Bangladesh	2004	Shastho Shebikas	25-35	F	150-300 HHS	PT	Educated
18.	(Campos et al., 2004)	Brazil	2004	CHA	-	M/F	150- 250 HHS	FT	Educated
19.	(Campos et al., 2004)	China	-	Bare foot doctor	-	-	--	-	-
20.	(____, 2005)	Egypt	2005	CHWs	-	75% M	-	FT	-
21.	(WHO, 2006c)	Papua New Guinea	2005	CHWs	-	-	-	FT	
	<b>Author</b>	<b>Country</b>	<b>Year</b>	<b>Name</b>	<b>Age</b>	<b>Gender</b>	<b>Coverage</b>	<b>Empl oy*</b>	<b>Level of Education</b>

22.	(Ismail B. & El, 2005)	Sudan	2005	CHWs	-	-	-	FT	-
23.	(Rosenthal, 2005)	USA	2005	CHW	30-39	F	-	75% Full Time	High School
24.	Myanmar(____, 2006)	Myanmar	2006	CHW	-	-	-	FT	Graduation
25.	(Keni, 2006)	Republic of Marshall Islands	2006	HA	26	M	-	FT	High School
26.	CAR(WHO, 2006d)	CAR	2006	CHW	-	-	--	FT	-
27.	Zimbabwe(WHO, 2006e)	Zimbabwe		VHW	-	-	1-3 villages	FT	--
28.	(Brown et al., 2006)	Peru	2006	CHW	19-70	M 75%	Varied	FT	Illiterate – Graduates
29.	(Sein, 2006 )	DPR Korea	1955	Sanitary Monitor	-	F	20-30 HHS	FT	-
30.		Myanmar	1976	CHW	-	F	200 HHS	FT	-
31.		Timor Leste	1978	Posyandu	-	F	10-20HHS	FT	-
32.		Indonesia	1978	Posyandau	-	F	10-220 HHS		
33.		USA	1960	CHW	30-50	F 80%	-	FT/PT	Graduates

#### CHWs in India

34.	Dave(Dave, 1991)	India/Maharastra (Sewagram)	1972	VHW	-	M			
35.	Dave(Dave, 1991)	India/West Bengal (CINI)	1975	CHW	-	F	-	-	
36.	Dave(Dave, 1991)	India/Gujarat (Tribhovandas Foundation)	1980	CHW	-	F			
37.	(Kumar, Deodhar, & Murthy, 1978)	India	1978	CHV	-	M/F	-	PT	-
38.	(Leslie, 1985) (Maru, 1983)	India	1977	CHV	>30	M/F	1/1000 ind	P/T 2-3 hrs	Primary schooling
39.	(Bhattacharji, Abraham, Muliyl, Job, John, & Joseph, 1986)	India/Vellore	1983	CHWs		F	1/1000 to 1500/ PTCHW	PT	Higher primary
40.	(UNICEF, 2004)	India	2000	VHGs		M	1/1000 HHS	PT	Literate
41.	(Mistry & Antia, 2003)	BANWASI*/ Uttar Pradesh/ India	2003			F	15/100 villages	PT	Primary education
42.		AWARE*/Andra Pradesh /India	2003			F	2/20 villages	PT	Primary education
43.		CINI*/ West Bengal / India	2003			F	1/400 families	PT	Primary education
44.		RUSHA*/ Vellore/ India	2003			F	1/1000 individuals	PT	Primary education
45.		FRCH*/Pune /India	2003	Gramsakhi		F	1/village	PT	Primary education

Abbreviations: \* Nature of employment; FT: Full Time, PT: Part Time; indi: Individual; VHW: Village Health Worker; HHS: Households; LHW: Lady Health Worker; RHA: Rural Health Assistant; CV: Community Volunteer; HA: Health Assistant; VHG: Village Health Guide  
Source: Compiled from Various Sources

**Table: 2. Management of CHWs under various programs**

Sl no.	Author	Count	Program	Training	Service provided	Monitor	Incentives
--------	--------	-------	---------	----------	------------------	---------	------------

		<b>ry</b>					
1.	(Hathirat, 1983)	Thailand	Abbots	3 weeks	PHC	-	Volunteer
2.	(Scholl, 1985)	Nicaragua	Brigadista	8 days	PHC, curative tasks	Health professionals	Volunteer/paid
3.	(Berman, 1984)	Java/Indonesia	CHD	Days/weeks	-	-	Volunteer
4.	(Bender & Pitkin, 1987)	Costa Rica	RCHP	16 wk	Updating census, immunization, treating malaria, health education, promoting FP, referral, participation in community organization	Physician	State government for training and supported by community
5.	(Bender & Pitkin, 1987)	Nicaragua	IOPAA	-	nutrition, sanitation, treat common disease, MCH care and Occupational Health	-	Voluntary, but report to the health system
6.	(Bender & Pitkin, 1987)	Colombia	-	-	First aid, child care, sanitation, treatment of common diseases Monthly visits to all households in the catchments area.	-	The resources were from the ministry of health, municipal and communities own resources
7.	(Robinson & Larsen, 1990)	Colombia	Colombia research national health care	3 Months	PHC	By auxiliary nurse	Rewards: salary from Health System
8.	(Reis et al., 1991)	Indonesia	Kader	-	GOBI, ORT	Health professionals	Paid by the system
9.	(Stekelenburg, Kyanamina, & Wolfers, 2003)	Kalabo/Zambia	PHC	-	PHC	-	Volunteer
10.	(Campos et al., 2004)	China	Barefoot Doctor	3-6 months	Primary health Care	MMT	Volunteers/ Kind
11.	(Campos et al., 2004)	Brazil	CHA	6-8 months	Health education, referrals	Municipal co-operation	Municipal Co-operation
12.	(UNICEF, 2004)	Bangladesh	BARC	21 days	PHC	POs	Profit by sale of drugs
13.	(UNICEF, 2004)	Bhutan	VHW	12 days	PHC	Block development committee	Voluntary
14.	(UNICEF, 2004)	Nepal	FCHV	15 days	PHC , ,	No supervision	Voluntary
15.	(____, 2005)	Egypt	CHWs	5-6 days	GOBI-FFF	TAHSEEN trainers	MOHP
16.	(Douthwaite & Ward, 2005)	Pakistan	LHWP	3 months	MCH service,FP, health promotion and education, first aids	-	MOH
17.	(Ismail B. & El, 2005)	Sudan	-	-	Comunty based heath service	-	Supported by the community
18.	(Melany, Ron, & Jane, 2006)	Ingham County/ USA	PITCH	-	Health insurance enrollment, smoking cessation,	-	Ingham county health department, Cost for the fiscal year 2005, \$ 252000
19.	(Whitley, Everhart, & Wright, 2006)	USA	-	-	Providing primary health care and health education	-	Volunteer
20.	(Perez et al., 2006)	USA	Community voices CHW Prog	2-3 months	Health insurance enrolment, Immunization, Asthma Management	-	Community voices organization (NGO)
21.	(US Department of Health and Human Services, 2007)	USA	CHW prog	On job	Member of delivery services,navigator,screeing and health education, out-reach enrolling informing agent and organizer for camps in community	Employer	Paid /Volunteer Employed, paid per hour \$ 13 to \$15
<b>Sl no</b>	<b>Author</b>	<b>Country</b>	<b>Program</b>	<b>Training</b>	<b>Service provided</b>	<b>Monitor</b>	<b>Incentives</b>
22.	(Kumar et al., 1978)	India	CHW scheme, 1978	6.6 weeks	PHC	-	Honorarium by government

23.	(Leslie, 1985) (Maru, 1983)	India	CHW scheme, 1977	3 months course Stipend 200/month	PHC	Voluntary workers from there village	Voluntary
24.	(Leslie, 1985)	India	CHW scheme INDIA		PHC	-	Rs. 200 during training, Rs. 50 per month
25.	(Bhattacharji et al., 1986)	India	Project/Vellore INDIA	20 days PTCHW  One year Health Aide	PHC	Two PTCHW by one Health Aide	-
26.	(Mistry & Antia, 2003)	India	NGO management of CHWs INDIA		PHC	-	FRCH- 100/worker
27.	(UNICEF, 2004)	India	VHG Scheme	3 months	PHC	Community	Voluntary

Abbreviations: RCHP: Rural Health Care Programme; IOPAA: Operational Integration from bottom ; PITCH: People Improving the Community Health; MMT: Mobile Medical Team; CHA: Community Health Assistant; MOH: Ministry of Health; LHWP: Lady Health Worker Program; CHD: Community Health Development; FCHV: Female Community Health Volunteer; PO: Program Officers, VHG: Village Health Guide

**Table 3 Summary of research articles showing health outcomes with introduction of CHWs**

SI no	Author	Country	Research questions/ Conceptual frame	Methodology	Results/issues
1.	(Zeighami, Zeighami, Javidian, & Zimmer, 1977)	Iran	To determine the health workers knowledge, attitude and practice about family planning and also to know the gender differences in effectiveness of family planning	A KAP survey was conducted after 14 months of training. The total samples of 1308 eligible couples were from two sites, project (658) and control site (650).	The health workers were able to double the usage of pills among the eligible couples and this was true for both sexes of health workers, maximum between the age groups 25 to 34 years.
2.	Bender E Deborah, Pitkin (Bender & Pitkin, 1987)	Costa Rica	The paper examined the evolution and current status of VHWs	An analysis of country's progress is done using sidels hypothesis of fundamental shift of wealth and power considering the PHC program	IMR 61.5/1000 in 1970 decreased to 19.1/1000 1980; U5 mortality decreased from 5.1/1000 in 1970 to 1.1/1000 in 1980
3.	Bender E Deborah, Pitkin(Bender & Pitkin, 1987)	Nicaragua	The paper examined the evolution and current status of VHWs	An analysis of country's progress is done using sidels hypothesis of fundamental shift of wealth and power considering the PHC program	Malaria decreased 39% from 1977-1983, polio eradicated, measles, whooping cough and tetanus extinct
4.	Bender E Deborah, pitkin (Bender & Pitkin, 1987)	Colombia	The paper examined the evolution and current status of VHWs	An analysis of country's progress is done using sidels hypothesis of fundamental shift of wealth and power considering the PHC program	1978-1982, extend basic service to 82% of popln. Polio vaccination 23% - 43%, DPT 22% - 37%, BCG 36% - 71% and measles 21%-50%
5.	(Chopra & Wilkinson, 1997)	Rural South Africa	Evaluate the immunization coverage among the rural south African children with use of CHW	study took place in Hlabisa health district of KwaZulu/Natal, South Africa, population of around 205,000 people. The programme has been running for 9 years, 1 CHW/100 households.	The immunization coverage was generally high. Immunization coverage was highest for all antigens in children who lived in areas with CHWs. There are no significant difference b/w two groups for BCG and measles coverage.
SI no	Author	Country	Research questions/ Conceptual frame	Methodology	Results/issues
6.	(Homer, Davis, & Brodie, 2000)	Australia	Evaluation/ St George Outreach Maternity Project (STOMP)	A randomized controlled trial was conducted with 1,089 women (550 in the experimental group and 539 in the control	STOMP group women also reported a higher perceived 'quality' of antenatal care compared with the control group. STOMP group women saw slightly more midwives and fewer doctors than control group women did.

7.	(Wayland, 2002)	Brazil	Evaluation of PACS program to improve PHC coverage. CHW regular performing their basic duties, health education and liaison b/w community and public health system	Data of Maternal and child health survey in Triunfo was used, that had a section designated to evaluate the performance of CHWs	35% of caregivers reported-CHWs visit previous month and 22% reported never been visited by a CHW, 34% reported they had never received hypochlorite solution, 49% never discussed their health problems with CHWs 45% discussed water treatment( major problem in the area) Sample of 180 households surveyed only 4 reported to have consulted CHW when their child fall ill.
8.	(Kovach & Worley, 2004)	Philadelphia/ USA	Relationship b/w CHWs and low income pregnant women	both qualitative and quantitative data; 1st focus group interviews 3 Mobile sites in north Philadelphia Self determination, decision-making ability, self-sufficiency were defined as empowerment Sample 168, in Phase I, 80 in Phase II	The mean self determination score postpartum, decision-making ability score postpartum, and self sufficiency score postpartum were significantly greater than their respective means at the time of program registration
9.	(Campos et al., 2004)	China and Brazil	Issues related to reorganization of CHWs, past present and future with two case studies	In depth case study analysis of barefoot doctors of China and Community health agents in Brazil	Barefoot doctor: CDR- 40/1000 in 50s came to 10/1000 in 1974, IMR 160/1000 in 50s came to 25/1000 in 1974
10.	(Jokhio, Winter, & Cheng, 2005)	Pakistan	--	Cluster Randomized control sampling of 7 subdistricts randomly assigned delivery kits to TBAs and LHWs. PHC outcome were preinatal and maternal mortality	The maternal deaths and prenatal deaths reduced in the intervention area. Referral to public health services was also encouraged, and correspondingly, a higher proportion of women in the intervention group than in the control group were referred to an emergency obstetrical care facility
11.	(Kotecha & Karkar, 2005)	India	Health status of integrated child development service workers	280 anganwadi workers AWW	- Anemia prevalence was 72.3% - Prevalence of severe, moderate and mild anemia among AWWs was 0.7%, 15.7% and 55.8% respectively the fundamental question of the capabilities of ICDS AWWs to provide for all the services and their capacity to imbibe from the training provided to them for NHED.
12.	(Delacollette, Stuyft, & Molima, 1996)	Katanga health zone Zaire	Evaluate the potential to reduce malaria morbidity and mortality	Quantitative, simple random sample of households	Increased in health seeking behavior CHWs- desired for further training and to be a part of health system. CHWs increased the workload of health care staff. Community expectations were higher, often dissatisfied with the limited service, least interested in contributing to the efforts of CHWs, administrative control over CHWs, no motivation by CHWs w.r.t community participation in Malaria control
13.	Schmeller Wilfried (Schmeller, 1998)	East Africa/ Dermatitis	The objective; were to determine the extent and severity of diseases in school and pre-school children in a rural community in western Kenya which includes treatment by trained CHWs	1993 & 1995 two separate epi survey, 40,000 popln- 13 primary schools, 5780 children from 4-16 years were examined for skin disease. Only typical cases were counted and were treated with 12 CHWs. The evaluation was done in 1995	Slight decrease in dermatoses b/w 1993 (32.4%), 1995(29.6%)., bacterial skin infections reduced from 12.7% to 10.8%. the most impressive change was a marked reduction in the extent and severity of skin diseases This study demonstrates that CHWs are able to deal successfully with the most important dermatoses in rural areas after a short training period
	<b>Author</b>	<b>Country</b>	<b>Research questions/ Conceptual frame</b>	<b>Methodology</b>	<b>Results/issues</b>

14.	Kelly M. Jane et.al (Kelly et al., 2001)	Kenya/ Childhood Illness	Ob: to characterize CHW performance using an algorithm for managing common childhood illness	3 cross sectional hospital based evaluation Observations of consultations using a check list CHW documentation of assessments findings, classification, and treatment for each sick children in standard form Repeat examination by clinician	Each CHW was evaluated with 1 or 2 OP /IP cases depending on the availability. 90% of CHWs made right diagnosis of malaria Many failed to identify symptoms, illness and administering right drugs Lack of regular supervision by professionals, continued education, complexity of the training modules led to poor performance.
15.	Akramul Islam(Islam, Wakai, Ishikawa, Chowdhury, & Vaughan, 2002)	Bangladesh	To compare the cost-effectiveness of the tuberculosis (TB) programme run by the Bangladesh Rural Advancement Committee (BRAC), which uses community health workers (CHWs), with that of the government TB programme which does not use CHWs.	TB statistics and cost data was collected from July 1996–June 1997 and cost per patient cured was calculated.	185 and 186 TB patients were treated by BRAC and government respectively. It was found that the cost per patient cured was US\$ 64 in the BRAC area compared to US\$ 96 in the government area. IT was also found that the BRAC and government TB control programmes appeared to achieve satisfactory cure rates using DOTS and the involvement of CHWs was found to be more cost-effective in rural Bangladesh.
16.	Joel D et. al. (Joel, Sathyaseelan, Jayakaran, Vijayakumar, Muthurathnam & Jacob, 2003)	South India	This study attempted to examine the knowledge of chronic psychosis among health workers of a rural community health program in South India.	Site: The Rural Unit for Health and Social Affairs (RUHSA), 80 CHWs volunteered to take part in the study. Avignette describing a typical patient with chronic psychosis was developed for the study	Seventy (87.5%) of subjects in the whole sample had at least one non-biomedical explanation for the psychosis (e.g. black magic, evil spirits as cause, non-disease concept, seeking treatment from traditional healers or temples and not seeking medical help).
17.	Ramos-Crequeira (Ramos-Crequeira, Torres, Crepaldi, Oliveira, Scazufca, Menezes et al., 2005)	Brazil	The aims of the present study were to apply and evaluate a simple and potentially cost-effective method of dementia case finding by community health workers (CHWs)	25 community health workers were trained to identify dementia cases in 2,222 people aged 65 and older in Piraju, a Brazilian town with 27,871 inhabitants.	CHWs identified 72 elderly people as being possible cases of dementia. Thus, 45 cases were confirmed according to the diagnostic examination, indicating a PPV of 62.5% for the procedure. The overall frequency of dementia was 2% in this population.
18.	(Leinberger-Jabari, 2005)	multinational	Review of 25 years of work in the community	The study included community-based organizations, hospitals and community clinics	CHWs were increasingly effective in providing outreach health care for population those were missed by the main stream. It was also found that CHWs were effective in providing health education and appropriate referrals for clients.
19.	Douthwaite Megan (Douthwaite & Ward, 2005)	Pakistan	To assess the impact of the LHW on the uptake of modern contraceptive methods	Interview with HHS and LHWs, complete profile of HHS was collected. A sample of 4277 currently married women in the LHW served areas	Higher levels of the use of contraceptives was seen in rural areas with LHWs
20.	(Bang et al., 2005)	Gadchiroli, India	Observation of cohort of neonates in preintervention of home-based neonatal care in rural gadchiroli.	Retrospective analysis of data from 39 villages compared between preintervention year 1995 to 1996 and intervention years 1996 to 2003	The low birth neonates declined from 11.3 to 4.7 % and preterm neonates by 33.3 to 10.2%, incidence of the sepsis, asphyxia, hypothermia and feeding problems, declined significantly; due to repeated visits made by village health workers (intervention periods) to houses educating mothers on hygiene, breast feeding, thermal care. Prevention and management of infections, management of neonatal sepsis with antibiotics, administration of Vitamin K injections by VHWS and refereeing cases to SEARCH hospital.
21.	CATALYST/TAHSEEN (____2005)	Egypt	Evaluation of outreach health workers	In depth interview of 816 outreach health workers was carried out.	Increase in knowledge of OC from 41% to 88%, breast lumps not as a result of menopause 48% to 95%, FGM 46% to 96%, Counseling of FP 48% to 91% --reference to local clinics
22.	S L Norris (Norris, Chowdhury, K. Van Le, Brownstein, Zhang, Jr et al., 2006)	Review	A systematic review was to examine the effectiveness of community health workers in supporting the care of persons with diabetes	Review was done using medical text words, CHWs, LHWs, volunteers, promoters and others in the electronic data bases especially in Medline till 2004	The 18 primary studies were published between 1986 and 2003 and included eight RCTs. Most of the studies were conducted in the USA. The majority of intervention participants were female (range 53–100%) and middle-aged. health care utilizations decreased in emergency

					visits by 38% and admissions by 53% and hospital admissions related to diabetes decreased from 25% in 1999 to 20% in 2002
23.	(Onwujekwe, Dike, Ojukwu, Uzochukwu, Ezumah, Shu et al., 2006)	South east Nigeria	Timelines of appropriate treatment for malaria with implementation of CHW	An intervention village (N=597 households) and non intervention village (N=600 households).	Pre and post intervention showed the preference of CHWs over self treatment at homes. The use of community health workers (CHWs) increased from 0% to 26.1% (p < 0.05), while self-treatment in the homes decreased from 9.4% to 0% (p < 0.05) after the implementation of the CHW strategy. Use of patent medicine dealers also decreased from 44.8% to 17.9% (p < 0.05) after CHW strategy was implemented.

**Table 4: Organizational Issues that influence CHWs performance**

SI no	Author	Country	Research Qs/Conceptual Frame	Methodology	Results/issues
1.	(Kumar et al., 1978)	India	- administrative response, to CHW scheme - community attitude and perception to CHW scheme; mainly on participation	Interview : 544 Officials, 203 village level workers, 299 CHWs, 6013 community members, 604 community leaders	1. fairness in selection of CHWs 2. training to CHWs were satisfactory for eg. CHWs scored 3 out 5 in malaria control tests 3. hurdles: non availability of medical officer, no stipend, non availability of manuals and lack of clarity by the government 4. gradual decline in the number of kits and drugs 5. majority of CHWs maintained records
2.	Hathirat Sant (Hathirat, 1983)	Thailand	Follow up evaluation of evaluation of the health care training for Buddhist abbots and ecclesiastical heads	A sample of 1600 Buddhist abbots and 400 ecclesiastical heads were selected and interviewed	- 82 % of Abbots and ecclesiastical had understood about primary health care; - 66 % provide health education - 57% Improve or educate nutrition. sanitation and environmental problems.,, - 75% Dispense modern drugs and 40 % dispensed herbal drugs - 29 % gave medical care
3.	Peter A Berman (Berman, 1984)	Indonesia	An evaluation of coverage and equity	Household survey of two sub-districts, Glagah and Beran	Coverage: 71 % of all children under five were weighed; 32% in beran and 39% in Glagah contacted VHWs for illness Equity: children under five in poorer community have above average probability of attending weighing sessions.
4.	Scholl (Scholl, 1985)	Nicaragua	An assessment of CHW in two sites	One urban and one rural site was selected, these were 2 PHCs among 33 which had brigadista working successfully according to standards set.	These brigadistas seem to be more a part of the professional health delivery team, than community-based workers who work semi autonomously and are accountable to the community first. It was also found that they were more dependent on auxiliary nurse midwives for directions.
5.	(Twumasi & Freund, 1985)	Zambia	Analyze the problems and issues arising with regards to community participation approaches to PHC	Theoretical issues through community participation research, literature review, and case study of CHWs	-1CHW/17 villages, no means of transport -Completely political issue of conflict b/w different actors and ways to tackle it.
6.	(Bhattacharji et al., 1986)	India	To evaluate the effectiveness of part time community health worker program	Sample 80,000 population	that educational status, experience, population covered, the degree of supervision and the scatter of houses all seem to influence performance. The age of the worker and the test scores do not seem to affect performance to a great extent. Supervision has an effect on Performance
SI no	Author	Country	Research Qs/Conceptual Frame	Methodology	Results/issues



7.	(Sauerborn, Nougara, & Diesfeld, 1989)	Burkina Faso	Recording utilization pattern of CHWs in a district of Burkina Faso	Household survey of N=715, HHS, 4 CHWs, 4 nurse midwives,	- 8.8 % CHWs detected mild disease - the villages bypassed CHWs 96.5% -no referral linkages b/w professionals and CHWs
8.	(Bentley, 1989)	Northwestern Somalia	Problem initiating a new health care program:CHWs	A case-control study; a village with a CHW service and with out Candidates – literate, preference female, b/w 20-40 yrs of age	Results revealed inadequate training, service bias, poor motivation. CHWs were satisfied with job and received kind from villagers for service but were not supported by health system.
9.	(Menon, 1991)	Gambia	Utilization of VHWs for PHC program	A HHS survey of mothers whose child had died last three years ; n=23	VHW provided preventive care. Mothers were not aware of VHW services and expected curative services a higher percentage of non-availability of VHWS reported
10.	Ruebush T. K. II, Weller, S. C., Klein R. (Ruebush et al., 1994)	Guatemala, malaria	The purpose of this investigation was to evaluate the criteria used by NMS workers to select volunteer community malaria workers and compare those criteria with the opinions of the residents about the qualities and characteristics they would prefer in an 'ideal' worker.	27 NMS, 7 sector chief and 100 residents of the Pacific Coast was selected. Interview as well as observational data was collected by spending half a day with 27 evaluators.  CVs = Community volunteers	11 qualities of an ideal CVs was brought out through open ended interview with households - takes care of pt at all times of the day even when busy -is at home all of the time -has general knowledge of medicine -is a responsible person - is interested in the welfare of his neighbors - recognizes the importance of his work as CVs - has the ability to learn the duties of a CV - is friendly -treats every one equally -is widely known in the community -is well liked
11.	(Curtale, Siwakoti, Lagrosa, LaRaja, & Guerra, 1995)	Nepal	the study tested the hypotheses that volunteers can provide effective PHC	One intervention and one control area, 2160 children total. Indepth interview with mothers of children was done to know the first contact with CHVs for the past 12 months. Along with a total of 208 CHVs were also include in the sample.	95% of mothers in the intervention met CHVs at least once compared to 24% in control group.35% mothers brought children to CHVs in intervention group. The ORS utilization was 78% in intervention group and 64% in the control group. The CHVs received double supervision and felt "not being" left alone.
12.	Roman and Lindsay (Roman et al., 1999)	Michigan/USA CISS program	Describe the perceptions of the benefits and stressors of helping as experienced by CHWs in an nurse-coordinated maternal child health intervention. Helpers Perception Measures, developed to assess benefits and stressors, were examined.	Part of Community Integrated service system program had two types of CHWs, paid and volunteers. Were given training to provide services to pregnant women who were at greater socio demographic and psychosocial risk than the staff had anticipated.	Highest ranking benefits included positive feelings associated with being involved in good work 95%, a sense of belonging 94% and greater self esteem 91%. They felt energized by helping others 81%. There are helper therapy benefits for CHWs who function in a maternal support program for low-income pregnant women.
13.	Ansari and Phillip (Ansari & Phillips, 2001)	South Africa	Aim: to compare the views of participants from four stakeholder groups as regards their voluntary status: the token-paid CHWs; the full-time employed projects' core staff ; unpaid 'solo' community members and, representatives of NGOs	self-administered questionnaire using snowballing' technique Benefits and costs of participation, satisfaction with partnership, sense of ownership, community representation, commitment and contribution  Out of 427 participants from various groups there were 70 CHWs	Benefits exceeded the costs General atmosphere of satisfaction Stakeholders and beneficiaries perceived a sense of ownership
<b>Sl no</b>	<b>Author</b>	<b>Country</b>	<b>Research Qs/Conceptual Frame</b>	<b>Methodology</b>	<b>Results/issues</b>
14.	(Dieleman, Cuong,	Vietnam	Develop strategies influencing staff	53 semi structured questionnaire was carried out, included 24 health staff	Motivating factors for health workers were appreciation by managers, colleagues and

	Anh, & Martineau, 2003)		motivation for better performance	and 6 receivers	the community, a stable job and income and training. The main discouraging factors were related to low salaries and difficult working conditions.
15.	(Lynn & Theresa, 2004)	---	Meta analysis 1971-1999, Difference b/w trained and untrained birth attendants on maternity care; KAP studies.	60 studies were included 44 developing countries, TBA assisted deliveries ranged from <1% to 66% live births	The results for TBA attributes were all positive, fairly uniform and significant The lowest estimate of 0.52 for 'behaviour' represents a 63% 'improvement' for trained TBAs over the untrained TBA baseline.
16.	(Brown et al., 2006)	Peru	Describing the profile of CHWs in Peru	Qualitative and quantitative research, community health projects from 1997- 2002, 40 andean communities, sample of 171 CHWs	CHWs more of young males, high school graduates, resulted in increase drop out rates, voluntary basis all these were in contract with traditional healers
17.	(Pahan, Prenger, Roy, & Pahan, 2007)	Bangladesh	To compare the advantages and disadvantages of local CHWs versus government practice contributing to improved service delivery for poor.	The study was conducted at the LAMB Integrated Rural Health and Development Project in North-West Bangladesh. 34 local CHWs compared with 11 externals. followed by 6 FGDs with community; in-depth interview with 17 representatives of two groups of CHWs	The community preferred local CHWs NGOs preferred more qualified external health worker than a less qualified internal health worker, for the simple reason that internal less qualified worker would reduce the performance of the NGOs.
18.	(Doherty & Coetzee, 2005)	South Africa	Relationship b/w CHWs and professional nurse	16 interviews and 1 FGD; with nurse and CHW. Age nurse 25-53; CHWs 30-55; predominantly women	nurses were unsure of the CHW role and CHW experienced being undermined initially. They were unaware of the training that CHWs had received. Nurses didn't accept CHWs because they were not professionally trained. CHWs wanted government to recognize them they felt as" not belonging CHWs began to understand the value of being in the community and nurses accept referrals from CHWs which was not the case earlier. Nurses stopped thinking CHWs as treat but as people who help them.
19.	(Sundaram an, 2005)	India	Why CHWs keep resurrected?, why and How NGOs have shown success?	Review of work of nine NGOs in India; who incorporated the concept of CHW	Success by NGO - good referral linkages -high quality leadership -women as health care providers -failure by Govt -male health workers -patronage, corrupted the choice -no continued training -weak referral -curative than preventive care
20.	(SARDI, n. d)	India	Working conditions, nature of work and targets, employment, job satisfaction, association with national and international allies	Coimbatore - 28, Chengalpet - 16, Madurai - 16; 54 VHNS; 6 MPWs	- health center located outskirts of villages -poor transport facilities -12 months to develop rapport with villages -lack of security -increase in the coverage area as the posts remain vacant -lack of financial incentives; difference b/w state & center -sexual harassment -over burdened with records entry -Urban health Posts: tasks unrelated to health department on workers -cover vacant posts -non- payment allowance -suspension on raising voice -both VHNS and MPws face enormous amount of mental stress

**Table 5. Financing CHW programs in Developed and Developing countries**

SI No	Author	Country	Research Qs/Conceptual Frame	Methodology	Results/issues
1.	Love Beth	San	This article will	Mail and telephonic survey in 8	65% of CHWs are full time and 35% are part

	Mary (Love, Gardner, & Legion)	Francisco	describe the functions and attributes of the Community Health Worker based on the findings of a systematic eight-county survey of the San Francisco Bay Area in 1996.	northern California counties with objectives: proportion of health care employers that employ CHWs, total no.of CHWs employed, demand for CHWs, profile of CHWs, barriers to wider employment. Out of 197 organisations in the region who responded for the survey, 71 (26%) either employ or plan to employ CHWs of which 62 were currently employing CHWs. A total of 504 CHWs are working in the 62 agencies reporting employing CHWs	time.44% of CHWs -fulltime salary - \$20,000 and \$25,000 93% of Agencies – provide health benefits 88%CHWs – government employees 66% organisations report to have career ladder 55%-salary-hard money (ongoing funding) 42% soft money (grants ,3yrs) Primary source- county/city funding 29%, federal grant 17%,66% CHWs are women –[African American 30%] 58% received formal level of education, 95% organization provided on job training; major conc. Of workers were from HIV/AIDS/STDs (27%), MCH (16%), alcohol and drugs 11%, primary care 10%, 91% - indicate budget constraints as a barrier to wider employment, 33% difficulty in supervising employers
2.	(Robinson & Larsen, 1990)	Colombia	Work Performance 'General Model of Work Behavior' The research was based on a theoretical model of worker performance that focuses on job related sources of rewards and feedback	The data are drawn from a broader study of health promoters (CHWs) . A survey research design was employed to obtain information from a random sample of rural health promoters (N = 179) and their auxiliary nurse supervisors about CHW performance and contributing factors	The findings indicate that feedback and rewards from the community have a greater influence on work performance. The findings do not support what appears to be a widely held assumption that the health system plays the primary role in influencing motivation and performance of CHWs
3.	(Thomason & Kolehmainen-Aitken, 1991)	Papua New Guinea	Performance of Rural Health staff ; identify the costs and the range of costs variation in health services and to assess outputs of rural health facilities	Survey was conducted among 76 rural health centers and 57 Churches	Inequitable unit cost of providing care was less than GO. Inequitable distribution on analysis with indicators for staffing need, more concentration on curative aspect. Church staff performed better than GO staff.
4.	(Makan & Bachmann, 1997)	South Africa	The aim of this study was to evaluate and analyse the nature, performance and costs of a sample of peri-urban and rural based CHW programs operating in the Western Cape province.	Three community based health care program were compared, 1517 households were interviewed in these areas, cost analysis of CHW program for the year 1994/95 fiscal year was done, compared with National Progressive Primary Health Care Network Training Centre (NPPHCN-TC)	The average cost for initial training at the NPPHCN-TC was approximately R17, 000 per CHW during 1994 and R10, 000 during 1995 and the average cost per visit to a CHW ranged from R11 to R35. For the three peri-urban CHW programs, the average cost per home visit was R26, R28, and R27 respectively. On an average the cost of visiting an out patient in a community health clinic is R55 and a normal clinic is R30. A patient visit to a CHW was generally less costly than a CHW home visit. CHWs average costs were less costly in the peri-urban areas than in rural areas.
5.	(Khan, Ahmed, & Saha, 2000)	Bangladesh	To estimate the additional time required for existing health worker to complete IMCI guidelines and also to estimate the number of new community health workers required for the same.	The data collection over a period of four months at two levels. One at the CHW level a sample of 1,921 cases and 3,584 cases at the paramedic level.	CHW took less than 20 min of time to examine 87 percent of children under IMCI guidelines. CHWs spent more time on diarrhea/ dysentery. With this an estimate of 4 to 6 hours per day was necessary for providing care. An estimate of 240 working hours per year would cost US\$ 992.
6.	(Ismail, S Immink, Mazar, & Nantel, 2003)	Kenya	Evaluation of CB-nutrition programm	Review on community food and nutrition program	low monetary incentive increase dropouts 50 families to be covered ZW\$ 500 (US\$ 10)/ CHW
7.	(Harter & Leier, n,d)	Canada	The impact of new economy on CHWs, income, work experience	Interview with N=836 Members of UFCW	- increase in stress due to job security followed by "Health and Social Services Delivery Improvement Act" - serve more people in less time, morale affected -reduction in no of trainings affected their performance -increased in incidents of injury -union views not addressed in new economy
<b>Sl No</b>	<b>Author</b>	<b>Country</b>	<b>Research Qs/Conceptual</b>	<b>Methodology</b>	<b>Results/issues</b>

			<b>Frame</b>		
8.	(Brown et al., 2006)	Peru	Describing the profile of CHWs	Describe the profile of CHW Qualitative and quantitative, 1997-2002 CH projects (41) n = 171	More young males with high school graduation Increase drop out rates among them, voluntary basis; completely opp was traditional healers
9.	Melany Mack, Ron Uken, Jane Powers (Melany et al., 2006)	Northwest Lansing, USA	Evaluation of enrolment of uninsured into Ingham Health Plan	Using three community-based organizations and Greater Lansing African American Health Institute, qualitative interview with the CHWs and quantitative data of Ingham Health Plan was collected	To start only 50 per cent of base line adults had coverage, with the introduction of CHWs, the enrolment increased substantially not only in Ingham health plan but also in Medicaid.
10.	Elizabeth M Whitley, Rachel M. Everhart, Richard A. Wright (Whitley et al., 2006)	Denver Health Community Voices, USA	The purpose of the study was to evaluate the financial effectiveness of CHW interventions with a population using a public safety net system; using return on investment way of cost analysis	A sample of all clients who began working with a CHW between January 1, 2003 and June 30, 2004 and had patient activity within the Denver Health system prior to their initial involvement with the CHW. Pre-intervention baseline data consisted of clients' utilization and charges that occurred during the 9 months before the initial intervention of a CHW.	Pre intervention cost - \$5,343,135, Post-intervention \$5,043,808 Increase in total visits from 5211 to 6630 , statistically significant was found in primary care.
11.	Moises Perez Sally E. Findley Miriam Mejia Jacqueline Martinez (Perez et al., 2006)	New York, USA	Evaluate the experiences of CHWs for health insurance, child immunizations, and asthma management from 2000-2005	"Descriptive and qualitative methods are used to demonstrate the extent and impact of the training programs on CHWs, the participating organizations, and community residents."	200% increase in insurance enrollment, 32% increase in asthma management, 16% immunization
12.	Sophie Witter(Witter, Kusi, & Aikins, 2007)	Ghana	Assess the impact of exemption of delivery fee scheme on health workers and TBAs	A cross sectional survey was done among the health workers, doctors, nurses, community health nurses and TBAs. The structured questionnaire was used to capture the household characteristics', income, working hours, and views about the exemption of the scheme.	The results showed that the professionals increased their working hours with relatively increase in workload to counterbalance their pay. The TBA suffered the most with the exemption of the scheme.

<b>Table 6: The SWOT analysis of CHW programs</b>	
<b>Strength</b>	<ul style="list-style-type: none"> <li>✓ CHWs are highly respected and valued in the communities by involving themselves in the community activities (Brown et al., 2006; Swider, 2002)</li> <li>✓ Community Based antenatal care approach has positive results(Homer et al., 2000)</li> <li>✓ LHWs are effective in providing modern contraceptives in rural areas. (Douthwaite &amp; Ward, 2005)</li> <li>✓ Empowerment of low income women (Kovach &amp; Worley, 2004)</li> <li>✓ CHW can be trained to perform wide range of PHC activities (Campos et al., 2004)</li> <li>✓ Promote equitable access to care (Berman, Gwatkin, &amp; Burger, 1987; Marguerite, Treadwell, &amp; Northridge, 2003)</li> <li>✓ Cost effective way of reaching underserved and inaccessible population (Walker &amp; Jan, 2005) (Andrews, Felton, Wewers, &amp; Heath, 2004; Berman et al., 1987)</li> <li>✓ CHWs are part of the community experience the same problems and can promote community organizations to confront the basic cause of ill health (Cruse, 1997)</li> <li>✓ Provide culturally appropriate health education and information by teaching concepts of health promotion and disease prevention (NRHA, 2000)</li> <li>✓ Highly accessible and highly trusted as CHW resides in the same village (Werner, 1977)</li> <li>✓ Low or no charges for service (Werner, 1977)</li> <li>✓ Highly effective than professionals in treating primary care (Werner, 1977)</li> </ul>
<b>Weakness</b>	<ul style="list-style-type: none"> <li>✓ More concentration on curative treatment (Thomason &amp; Kolehmainen-Aitken, 1991)</li> <li>✓ CHW selection, not known to community, lack of logistic support, lack of incentives to maintain records, no incentive for working(Stekelenburg et al., 2003)</li> <li>✓ Complexity of guidelines for management of sick children(Kelly et al., 2001)</li> <li>✓ Non- standardization / certification of CHW education (Doherty &amp; Coetzee, 2005)</li> <li>✓ Not recognized as legitimate providers(Assembly, 2006)</li> <li>✓ Absenteeism, poor quality of work, low morale, weak organizational and managerial issue; have resulted in lower performance of CHW(Berman, 1984; Berman et al., 1987; McElmurry, Marks, Cianelli, &amp; Mamede, 2002)</li> <li>✓ Low community participation, villagers not involved in identification of problems and lesser duration of training lowered the performance (Sauerborn et al., 1989)</li> <li>✓ lack of definite work schedule (Sringernyuang et al., 1995)</li> <li>✓ Programs must be adequately funded (Cruse, 1997)</li> <li>✓ Lack of logistic support (Zuvekas, Nolan, &amp; Tumaylle, 1998)</li> </ul>
<b>Opportunities</b>	<ul style="list-style-type: none"> <li>✓ CHWs are becoming increasingly effective members of the health care delivery team because they are able to provide outreach services to communities been missed through larger main stream organization (Leinberger-Jabari, 2005)</li> </ul>

	<ul style="list-style-type: none"> <li>✓ Rewards from community have a direct effect on performance (Robinson &amp; Larsen, 1990)</li> <li>✓ Educating and motivating women to receive antennal care showed increased utilization of health facility (Lynn &amp; Theresa, 2004)</li> <li>✓ Integrating TBAs and LHWs with health care system would reduce perinatal mortality and maternal deaths.(Jokhio et al., 2005)</li> <li>✓ CHWs gained valuable work experience (Roman et al., 1999)</li> <li>✓ Increased under five immunization coverage(Chopra &amp; Wilkinson, 1997)</li> <li>✓ Increased utilization of health facility and enrollment into health insurance (Assembly, 2006)</li> <li>✓ Act as a two way referral mechanism between community and the professionals at the health system. (Marguerite et al., 2003)</li> </ul>
<b>Threats</b>	<ul style="list-style-type: none"> <li>✓ Inadequate training , service bias and poor motivation would lead to lower levels of confidence among CHWs (Bentley, 1989)</li> <li>✓ Politicization of conflict issues between different providers would hamper the role of CHWs to meet objectives (Twumasi &amp; Freund, 1985; Zuvekas et al., 1998)</li> <li>✓ Lower levels of trust in CHWs and lack of intersectoral collaboration will lead to bypassing CHWs for referrals (Sauerborn et al., 1989) (Cruse, 1997)</li> <li>✓ If the felt needs of the community are not addressed by the programs (Wayland, 2002)</li> <li>✓ Non-financial incentives not accounted as a motivating factor for performance by CHWs (Dieleman et al., 2003)</li> <li>✓ Lack of government policies, poor interpersonal relation with the government health staff, community and professionals, lack of supervision and continued support, will add to poor performance (Campos et al., 2004; Gilson, Walt, Heggenhougen, Owuor-Omondi, Perera, Ross et al., 1989)</li> <li>✓ Lack of defined roles and responsibilities of health workers in relation with CHWs (Assembly, 2006; Zuvekas et al., 1998)</li> <li>✓ If CHW observed as a part of publicly funded health system, they lose the instinct to serve the community (Anne &amp; Taati, 2003)</li> <li>✓ No existing functional health infrastructure hampers referrals (Bentley, 1989; Zuvekas et al., 1998)</li> <li>✓ To work continuously as CHW with out expecting any change in designation (Sringernyuang et al., 1995)</li> <li>✓ Willingness of the community to retain CHW scheme (Sringernyuang et al., 1995)</li> <li>✓ Considering monitory incentive as “salary” would increase drop out rates (S. Ismail et al., 2003)</li> <li>✓ When CHWs is seen as cheap substitutes to the regular health staff leads to death of the program (Cruse, 1997)</li> <li>✓ Low patient demand and competing interest result in attrition(Gray &amp; Ciroma, 1988)</li> </ul>

## Reference:

- \_\_\_\_\_, (2005). Best Practices in Egypt: Mobilizing Community out reach workers. State-of-the-art Reproductive Health and Family Planning Services: The Catalyst Consortium/TAHSEEN.
- \_\_\_\_\_, (2006). Government and Policy: Ministry of Health. Myanmar Online Data Information Network Solutions, retrieved from [www.modins.net/MyanmarInfo/ministry/health.htm](http://www.modins.net/MyanmarInfo/ministry/health.htm), accessed on 9<sup>th</sup> February, 2007
- Abbatt, F. (2005). Scaling up Health and Education Workers: Community Health Workers. London: DFID Health Systems Resource Centre.
- Ande, O., Oladepo, O., & Brieger, W. R. (2004). Comparison of Knowledge on diarrheal disease management between two types of community based distributors in Oyo State Nigeria. . *Health Education and Research*, 19(1), 110-113.
- Andrews, J. O., Felton, G., Wewers, M. E., & Heath, J. (2004). Use of Community Health Workers in research with ethnic minority women. *Journal of Nursing Scholarship*, 36(4), 358-365
- Anne, L., & Taati, I. (2003). Adding Value and equity to primary health care through partnership working to establish a viable community health workers' programme in Namibia. *Critical Public Health*, 13(4), 331-346.
- Antia, N. H., & Bhatia, K. (1993). NGO Experiences: Implications for health services at the community level. In FRCH (Ed.), *People's health in people's hands*. Mumbai: Foundation for Research in Community Health.
- Ansari, W. E., & Phillips, C. J. (2001). Inter professional Collaboration: a stakeholder approach to evaluation of voluntary participation in community partnerships. *Journal of Inter professional Care*, 15(4), 351 - 368
- Arole, S. (2007). Community health workers: the experience of CRPH, Jamkhed. *Health Action For All*, 20(4).
- Ballester, G. (2005). Community Health Workers: Essential to Improving Health in Massachusetts. Bureau of Family and Community Health. Boston: Massachusetts Department of Public Health, retrieved from [www.mass.gov/dph](http://www.mass.gov/dph)
- Bang, A. T., Bang, R. A., P. G., Sontakke & the SEARCH team. (1994). Management of Childhood pneumonia by traditional birth attendants. *Bulletin of the World Health Organisation*, 72(6), 897-905.

- Bang, A.T., Baitule, S. B., H. M., Reddy, M. D., Deshmukh, R. A., Bang (2005). Low Birth Weight and Preterm Neonates: can they be managed at home by mother and a trained village health worker? *Journal of Perinatology*, 25:S72-S81
- Bender, D. E., & Pitkin, K. (1987). Bridging the gap: The village health worker as the cornerstone of the primary health care model. *Social Science & Medicine*, 24(6), 515-528.
- Bentley, C. (1989). Primary Health Care in Northwestern Somalia: A case Study *Social Science Medicine*, 28(10), 1019 - 1030.
- Berman, P. A. (1984). Village health workers in Java, Indonesia: Coverage and equity. *Social Science & Medicine*, 19(4), 411-422.
- Berman, P. A., Gwatkin, D. R., & Burger, S. E. (1987). Community-based health workers: Head start or false start towards health for all? *Social Science & Medicine*, 25(5), 443-459.
- Bhattacharji, S., Abraham, S., Muliylil, J. P. P., Job, J. S., John, K., & Joseph, A. (1986). Evaluating Community health workers performance in India. *Health Policy and Planning*, 1(3), 232-239.
- Bhattacharyya, K., Winch, P., LeBan, K., & Tien, M. (2001). Community Health Worker incentives and Disincentives: How they affect motivation, retention, and sustainability, Basic support for Institutionalizing Child Survival Project (BASICS II). Arlington, Virginia. : The United States Agency for International Development. Retrieved from [www.basics.org](http://www.basics.org)
- Brown, A., Malca, R., Zumaran, A., & Miranda, J. J. (2006). On the front line of primary health care: the profile of community health workers in rural Quechua communities in Peru. (p. 11). Retrieved from [www.human-resource-health.com/content/4/1/11](http://www.human-resource-health.com/content/4/1/11)
- Campos, F. E. d., Ferreira, J. R., Souza, M. F. d., & Aguiar, R. A. T. d. (2004). Innovations in Human Resources Development: the Role of Community Health Workers. Joint Learning Initiative- working group 6.
- Chopra, M., & Wilkinson, D. (1997). Vaccination Coverage is higher in Children Living in Areas with Community Health Workers in Rural South Africa. pp. 372-374). Retrieved from <http://tropej.oxfordjournals.org>
- Couper, I. (2004). Rural Primary Health care in Iran. *The South African Academy of Family Practice*, 46(5). Retrieved from [www.safpj.co.za/index.php/safpj/article/viewFile/81/81](http://www.safpj.co.za/index.php/safpj/article/viewFile/81/81)



- Cruse, D. (1997). Community Health Workers in South Africa: information for provincial policy makers. Durban: The Health Systems Trust, available on <ftp://ftp.hst.org.za/pubs/other/chws.pdf>
- Curtale, F., Siwakoti, B., Lagrosa, C., LaRaja, M., & Guerra, R. (1995). Improving skills and utilization of Community Health Volunteers in Nepal. *Social Science & Medicine*, 40(8), 1117-1125.
- Dave, P. (1991). Community and self-financing in voluntary health programmes in India. *Health Policy and Planning*, 6(1), 20-31.
- Delacollette, C., Stuyft, P. V. d., & Molima, K. (1996). Using Community Health Workers for malaria control: Experience in Zaire. *Bulletin of the World Health Organisation*, 74(4), 423-430.
- Dieleman, M., Cuong, P., Anh, L., & Martineau, T. (2003). Identifying factors for job motivation of rural health workers in North Viet Nam. (p. 10). Retrieved from [www.human-resource-health.com/content/1/1/10](http://www.human-resource-health.com/content/1/1/10)
- Doherty, T. M., & Coetzee, M. (2005). Community Health Workers and Professional Nurses: Defining the Roles and Understanding the Relationships. *Public Health Nursing*.22:360-365. available from [www.blackwell-synergy.com/doi/abs/10.1111/j.0737-1209.2005.220413.x](http://www.blackwell-synergy.com/doi/abs/10.1111/j.0737-1209.2005.220413.x)
- Douthwaite, M., & Ward, P. (2005). Increasing contraceptive use in rural Pakistan: an evaluation of the Lady Health Worker Programme. *Health Policy and Planning*, 20(2), 117-123.
- Friedman, I. (2005). Community Health Workers and Community caregivers towards unified model of practice: Health Systems Trust.
- Gilson, L., Walt, G., Heggenhougen, K., Owuor-Omondi, L., Perera, M., Ross, D., et al. (1989). National Community Health Worker Programs: How can they be strengthened? *Journal of Public Health Policy*, 10(4), 518-532.
- GOI. (1956). The second five year plan (1956-1961), The Five Year Plans. The Planning Commission, chapter 25, New Delhi.
- GOI. (1997). Performance Budget 1995- 96, In MOHFW (Ed.), Performance Budgets. Chapter 17; New Delhi: Ministry of Health and Family Welfare. Retrieved from <http://mohfw.nic.in/reports/Performance%20Budget1995-96/part2/Ch17.pdf>
- Gray, H. H., & Ciroma, J. (1988). Reducing attrition among village health workers in rural Nigeria. *Socio-Economic Planning Sciences*, 22(1), 39-43.

- Gryboski, K., Yinger, N. V., Dios, R. H., Worley, H., & Fikree, F. F. (2006). Working With the Community for Improved Health. Health Bulletin Washington, DC: Population Reference Bureau.
- Haines, A., Sanders, D., Lehmann, U., Rowe, A. K., Lawn, J. E., Jan, S., et al. (2007). Achieving child survival goals: potential contribution of community health workers. *The lancet*. 369 (9579), 2121-2131
- Harter, J.-H., & Leier, M. (n,d). No Time to Care: Community health workers in the New Economy. Center for Labour Studies at Simon, Fraser University. Retrieved from [www.sfu.ca/labour/UFCW-healthcare-study-Oct19.pdf](http://www.sfu.ca/labour/UFCW-healthcare-study-Oct19.pdf) , accessed on April 15, 2007
- Hathirat, S. (1983). Buddhist monks as community health workers in Thailand. *Social Science & Medicine*, 17(19), 1485-1487.
- Homer, C. S. E., Davis, G. K., & Brodie, P. M. (2000). What do women feel about community-based antenatal care? *Australian and New Zealand Journal of Public Health*, 24(6), 590 - 595.
- Islam, M. A., Wakai, S., Ishikawa, N., Chowdhury, A. M. R., & Vaughan, J. P. (2002). Cost-effectiveness of community health workers in tuberculosis control in Bangladesh. *Bulletin of the World Health Organization*(80), 445-450.
- Ismail, B., & El, S. B. (2005). Mapping of Human Resource for Health, Sudan Final Report. In F. Sudan (Ed.).
- Ismail, S., Immink, M., Mazar, I., & Nantel, G. (2003). Community - Based food and nutrition programmes: what makes them successful. A review and analysis of experience (p. Annex 2 (C)). Rome: Food and Agriculture Organisation of the United Nations.
- Jerden, L., Hillervik, C., Hansson, A.-C., Flacking, R., & Weinehall, L. (2006). Experiences of Swedish community health nurses working with health promotion and a patient-held health record. pp. 448-454).
- JLI. (2004). Human Resource for Health: Overcoming the crisis Global Health Trust.
- Jobert, B. (1985). Populism and Health Policy: The case of Community Health Volunteers in India. *Social Science Medicine*, 20(1), 1-28.
- Joel, D., Sathyaseelan, M., Jayakaran, R., Vijayakumar, C., Muthurathnam, S., & Jacob, K. S. (2003). Explanatory models of psychosis among community health workers in South India. *Acta Psychiatrica Scandinavica*, 108 (1), 66-69.

- Jokhio, A. H., Winter, H. R., & Cheng, K. K. (2005). An Intervention Involving Traditional Birth Attendants and Perinatal and Maternal Mortality in Pakistan. pp. 2091-2099).
- Kallander, K., Tomson, G., Nsabagasani, X., Sabiiti, J. N., Pariyo, G., & Peterson, S. (2006). Can community health workers and caretakers recognize pneumonia in children? Experiences from western Uganda. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 100(10), 956-963.
- Kash, B. A., May, M. L., & Tai-Seale, M. (2007). Community health worker training and certification programs in the United States: Findings from a national survey. *Health Policy*, 80(1), 32-42.
- Kelly, J. M., Osamba, B., & Grag, R. M. (2001). Community Health worker performance in the management of Multiple Childhood illness: Siaya district, Kenya 1997-2001. *American Journal of Public Health*, 91, 1617 - 1624.
- Keni, B. (2006). Training competent and effective Primary Health Care Workers to fill a void in the outer islands health service delivery of the Marshall Islands of Micronesia. (p. 27).
- Khan, M. M., Ahmed, S., & Saha, K. K. (2000). Implementing IMCI in a Developing Country: Estimating the Need for Additional Health Workers in Bangladesh. *Human Resources for Health Development Journal*, 4(2).
- Kotecha, P. V., & Karkar, P. D. (2005). How Anemic are ICDS Anganwadi Workers ? *Indian Journal of Community Medicine*, 30(4).
- Kovach, A. C., & Worley, J. B. H. (2004). The impact of community health workers on the self-determination, self-sufficiency, and decision-making ability of low-income women and mothers of young children. pp. (343-356).
- Kumar, S., Deodhar, N. S., & Murthy, N. (1978). An Evaluation of Community Health Workers Scheme - A Collaborative Study (1978) In S. a. Demography (Ed.): Central Council of Health and Family Welfare.
- Lehmann, U., Friedman, I., & Sanders, D. (2004). Review of the Utilization and Effectiveness of Community-Based Health Workers in Africa. *JLI WORKING PAPER 4-1: HUMAN RESOURCES FOR HEALTH AND DEVELOPMENT*.
- Leinberger-Jabari, A. (2005). VOICES OF THE COMMUNITY:A Profile of Minnesota's Community Health Workers. Voices of the Community: MIHV Publication.
- Leslie, C. (1985). What caused India's massive community health workers scheme: A sociology of knowledge. *Social Science & Medicine*, 21(8), 923-930.

- Lewin, S., Dick, J., Pond, P., Zwarenstein, M., Aja, G., Wyk, B. v., et al. (2005). Lay health workers in primary and community health care. The Cochrane Database of Systematic Reviews: John Wiley & Sons, Ltd.
- Love, M. B., Gardner, K., & Legion, V. Community Health Workers: Who Are They and What Do They Do? : Department of Health Education and City College of San Francisco,.
- Low, A., & Ithindi, T. (2003). Adding value and equity to primary healthcare through partnership working to establish a viable community health workers' programme in Namibia. *Critical Public Health*, 13(4), 331-346.
- Low, L. K., Moffat, A., & Brennan, P. (2006). Doulas as Community Health Workers: Lessons Learned from a Volunteer Program. *Journal of Perinatal Education*, 15(3), 25-33.
- Lynn, S., & Theresa, S. A. (2004). What can meta analysis tell us about traditional birth attendants training and pregnancy outcomes? . *Midwifery*, 20, 51-60.
- Magongo, B. (2004). Community Health Worker in Gautang: Context and Policy. The Gauteng Department of Health.
- Makan, B., & Bachmann, M. (1997). An Economic analysis of community health worker programmes in the western Cape province. In H. S. Trust (Ed.). Durban.
- Management, O. P. (2002). Lady Health Worker Programme: External Evaluation of the National Programme for Family Planning and Primary Health Care 1999-2002. Training Programme Review: Oxford Policy Management.
- Marguerite, R. J., Treadwell, H. M., & Northridge, M. (2003). Promoting Good Health. A Community Voices Publication National Center for Primary Care.
- Maru, R. M. (1983). The Community Health Volunteer Scheme in India: An Evaluation. *Social Science & Medicine*, 17(19), 1477-1483.
- Matomora, M. K. S. (1989). Mass produced village health workers and the promise of primary health care. *Social Science & Medicine*, 28(10), 1081-1084.
- May, M. L., & Contreras, R. B. (2006). Promotor(a)s, the organizations in which they work, and an emerging paradox: How organizational structure and scope impact promotor(a)s' work. *Health Policy*.
- Mburu, F. M. (1994). Whither community health workers in the age of structural adjustment? *Social Science & Medicine*, 39(7), 883-885.
- McElmurry, B. J., Marks, B. A., Cianelli, R., & Mamede, M. (2002). PRIMARY HEALTH CARE IN THE AMERICAS: CONCEPTUAL FRAMEWORK,

EXPERIENCES, CHALLENGES AND PERSPECTIVES. *Pan American Health Organization.*

- Melany, M., Ron, U., & Jane, P. (2006). People Improving the Community's Health: Community Health Workers as Agents of Change. *Journal of Health Care for the Poor and Underserved*, 17, 16-25.
- Menon, A. (1991). Utilization of village health workers with in a primary health care programme in the Gambia. *Journal of Tropical Medicine and Hygiene*, 94(4), 268-271.
- Mistry, N., & Antia, N. (2003). Community Based Health Workers- A review from India. The Foundation for Research in Community Health. Mumbai.
- National Human Services Assembly (2006). Community Health Workers: Closing Gaps in Families' Health Resources. Family Strengthening Policy Center, retrieved from [www.nassembly.org/fspc](http://www.nassembly.org/fspc)
- Norris, S. L., Chowdhury, F. M., K. Van Le, T. H., Brownstein, J. N., Zhang, X., Jr, L. J., et al. (2006). Effectiveness of Community Health Workers in the care of persons with diabetes. *UK Diabetic Medicine*, 23(5), 544-556.
- NRHA. (2000). COMMUNITY HEALTH ADVISOR PROGRAMS. In N. R. H. Association (Ed.), NRHA Rural Health Policy Board
- Nyonator, F. K., Awoonor-Williams, J. K., Phillips, J. F., Jones, T. C., & Miller, R. A. (2005). The Ghana Community-based Health Planning and Services Initiative for scaling up service delivery innovation. (pp. 25-34).
- Ofosu-Amaah, V. (1983). National Experience in the use of Community Health Workers. Geneva: WHO.
- Onwujekwe, O., Dike, N., Ojukwu, J., Uzochukwu, B., Ezumah, N., Shu, E., et al. (2006). Consumers stated and revealed preferences for community health workers and other strategies for the provision of timely and appropriate treatment of malaria in southeast Nigeria. (p. 117).
- Pahan, S., Prenger, K. L., Roy, J. C., & Pahan, F. (2007). Comparative Advantages of Local Community Health Workers Over External Professionals. Annual Scientific Conference (ASCON) 2007 Dhaka, Bangladesh: ICDDR,B retrieved from <http://www.icddr.org/pub/publication.jsp?classificationID=1&pibID=8120>
- Perez, M., Findley, S. E., Mejia, M., & Martinez, J. (2006). The Impact of Community Health Worker Training and Programs in New York City. *Journal of Health Care for the Poor and Underserved* 17, 26-43.

- Ramos-Crequeira, A. T. A., Torres, A. R., Crepaldi, A. L., Oliveira, N. I. L., Scazufca, M., Menezes, P. R., et al. (2005). Identification of Dementia Cases in the Community: A Brazilian Experience. *Journal of American Geriatric Society*, 53, 1738-1742.
- Ramprasad, V. (1988). Community health workers-an evolving force. *World Health Form*, 9, 239-234.
- Reis, T., Elder, J., Satoto, kodyat, B., & Plamer, A. (1991). An examination of the performance and motivation of Indonesian village health volunteers. *International Quaterly of Community Health Education (Abstract)*, 11(1), 19-27.
- Robinson, S. A., & Larsen, D. E. (1990). The relative influence of the community and the health system on work performance: A case study of community health workers in Colombia. *Social Science & Medicine*, 30(10), 1041-1048.
- Roman, L. A., Lindsay, J. K., Moore, J. S., & Shoemaker, A. L. (1999). Community Health Workers: Examining the Helper Therapy Principle. *Public Health Nursing*, 16(2), 87-95.
- Rosenthal, E. L. (2005). Program Assistance Document for the Community Health Worker Field. Migrant Health Promotion.
- Ruebush, T. K., Weller, S. C., & Klein, R. E. (1994). Qualities of an ideal volunteer community malaria worker: A comparison of the opinions of community residents and national malaria service staff. *Social Science & Medicine*, 39(1), 123-131.
- Salmen, L. F. (2002). Beneficiary Assment, an approach described. In E. a. S. S. Development (Ed.), *Social Development series Washington D C World Bank*.
- SARDI. (n. d). Health Workers- Life, Work and Collectivism. Labour: Mobility and HIV. Available from [www.mobilityandhiv.org/docs/Edited%205.doc](http://www.mobilityandhiv.org/docs/Edited%205.doc), accessed on November 26<sup>th</sup> 2006.
- Sauerborn, R., Nougara, A., & Diesfeld, H. J. (1989). Low utilization of community health workers: Results from a household interview survey in Burkina Faso. *Social Science & Medicine*, 29(10), 1163-1174.
- Schmeller, W. (1998). Community health workers reduce skin diseases in East African children. *International Journal of Dermatology*, 37, 370-377.
- Scholl, E. A. (1985). An assessment of community health workers in Nicaragua. *Social Science & Medicine*, 20(3), 207-214.
- Scott, G., & Wilson, R. (2006). Community Health Worker Advancement: A Research Summary. In R. W. J. Foundation (Ed.), *Jobs for the Future*. Boston: Skill works.

- Sein, U. T. (2006 ). Health Volunteers: Third Workforce for Health-for-All Movement. *Regional Health Forum*, vol 10(no 1), 38 - 48.
- Sringernyuang, L., Hongvivatana, T., & Pradabmuk, P. (1995). Implications of Community Health Workers distributing drugs: a case study of Thailand. Action Programme on Essential Drugs. England: World Health Organization.
- Stekelenburg, J., Kyanamina, S. S., & Wolffers, I. (2003). Poor performance of community health workers in Kalabo District, Zambia. *Health Policy*, 65(2), 109-118.
- Stock-Iwamoto, C., & Korte, R. (1993). Primary health workers in North East Brazil. *Social Science & Medicine*, 36(6), 775-782.
- Sundararaman, T. (2005). Review of Health Care in India. In L. V. Gangolli, R. Duggal & A. Shukla (Eds.), *Community Health Worker Programs and the Public Health Systems: Centre for Enquiry into Health and Allied Themes*.
- Sundararaman, T. (2007). Community Health Workers: scaling up programmes. *The Lancet*. 369(9579), 2058-2059
- Swider, S. M. (2002). Outcome Effectiveness of Community Health Workers: an integrative Literature Review *Public Health Nursing*, 19(1), 11-20.
- Thomason, J. A., & Kolehmainen-Aitken, R.-L. (1991). Distribution and performance of rural health workers in Papua New Guinea. *Social Science & Medicine*, 32(2), 159-165.
- Trust, G. H. (2003). Specific Programs and Human Resources: Addressing a key implementation constraint Human resources constraints in health services delivery: Global Health Trust.
- Twumasi, P. A., & Freund, P. J. (1985). Local politicization of primary health care as an instrument for development: A case study of community health workers in Zambia. *Social Science & Medicine*, 20(10), 1073-1080.
- UNICEF. (2004). What Works for Children in South Asia :COMMUNITY HEALTH WORKERS. In UNICEF, South Asia (Ed.). Katmandu, Nepal.
- US Department of Health and Human Services. (2007). Community Health Worker National Workforce Study. In B. o. H. Professions (Ed.), *The Community Health Workforce*. George Washington: Health Resource and Service Administration.

- Walker, D. G., & Jan, S. (2005). How do we determine whether community health workers are cost-effective?some core methodological issues. *Journal of Community Health*, 30(3).
- Wayland, C. (2002). Acceptable and appropriate: program priorities vs. felt needs in a CHW program. *Critical Public Health*, 12(4).
- Werner, D. (1977). THE VILLAGE HEALTH WORKER --LACKEY OR LIBERATOR? : Health Wrights Organization.
- Whitley, E. M., Everhart, R. M., & Wright, R. A. (2006). Measuring Return on Investment of Outreach by Community Health Workers. *Journal of Health Care for the Poor and Underserved*, 17, 6-15.
- WHO. (1989). Strengthening the performance of community health workers in primary health care. Technical Report Series 780. Geneva: WHO.
- WHO. (2006a). The World Health Report 2006- Working together for health. The World Health Report. New Delhi: WHO.
- WHO. (2006b). Country Health System Fact Sheet 2006:Algeria. WHO.
- WHO. (2006c). Health Situation in Papua New Guinea. In WPRO.WHO (Ed.), COUNTRY HEALTH INFORMATION PROFILES: WHO.
- WHO. (2006d). Country Health System Fact Sheet 2006:Central African Republic. WHO.
- WHO. (2006e). Country Health System Fact Sheet 2006:Zimbabwe. WHO.
- Witter, S., Kusi, A., & Aikins, M. (2007). Working practices and incomes of health workers: evidence from an evaluation of a delivery fee exemption scheme in Ghana. *Human Resources for Health* 5(2).
- Zeighami, E., Zeighami, B., Javidian, I., & Zimmer, S. (1977). The Rural Health Worker as a Family Planning Provider: A Village Trial in Iran. *Studies in Family Planning*, 8(7), 184-187.
- Zuvekas, A., Nolan, L. S., & Tumaylle, C. (1998). Impact of Community Health Workers on access, use of services and patient knowledge and behavior. In U. P. H. S. The Bureau of Primary Health Care (Ed.). George Washington: Center for Health Policy Research.